

CONSTRUCTION OF STRING SOLUTIONS AROUND NON-TRIVIAL BACKGROUNDS

C. Kounnas *

Theoretical Physics Division, CERN
CH - 1211 Geneva 23

ABSTRACT

We present a way of constructing string solutions around non-trivial gravitational backgrounds. The proposed solutions are constructed using $N = 4$ superconformal building blocks with $\hat{c} = 4$. We give two different and inequivalent realizations of non-trivial four-dimensional subspaces, and we show the emergence of the $N = 4$ globally defined superconformal symmetry. The existence of $N = 4$ world-sheet symmetry stabilizes our solutions and implies in target space a number of covariantized supersymmetries around space-time dependent gravitational and dilaton backgrounds.

*Talk given at the
International Workshop on
“String Theory, Quantum Gravity and Unification of Fundamental Interactions”
Rome, 21-26 September 1992
and at the
26th Workshop: “From Superstrings to Supergravity”
Erice - Sicily, 5-12 December 1992*

CERN-TH.6790/93

January 1993

*On leave from Lab. de Physique Théorique de l' ENS, F-75231, Paris, FRANCE.

CONSTRUCTION OF STRING SOLUTIONS AROUND NON-TRIVIAL BACKGROUNDS

Costas KOUNNAS[†]

Theoretical Physics Division, CERN
CH - 1211 Geneva 23

ABSTRACT

We present a way of constructing string solutions around non-trivial gravitational backgrounds. The proposed solutions are constructed using $N = 4$ superconformal building blocks with $\hat{c} = 4$. We give two different and inequivalent realizations of non-trivial four-dimensional subspaces, and we show the emergence of the $N = 4$ globally defined superconformal symmetry. The existence of $N = 4$ world-sheet symmetry stabilizes our solutions and implies in target space a number of covariantized supersymmetries around space-time dependent gravitational and dilaton backgrounds.

A way to better understand string theories and their induced low-energy field theories coupled to Einstein gravity is to study classical string solutions in the presence of non-trivial space-time backgrounds.

There are two ways to proceed in this direction:

- (i) The first is to use a two-dimensional σ -model where the non-trivial backgrounds correspond to some two-dimensional field-dependent coupling constants. The vanishing of the corresponding β -functions is then identified with the background field equation of motion in the target space-time [1].
- (ii) The second approach consists of constructing directly some underlying non-trivial conformal field theory, and then trying to interpret the obtained string vacuum in terms of the target space-time backgrounds.

The two methods are useful and complementary. The σ -mode approach provides a clear geometric interpretation, but it has the disadvantage of the α' -expansion which is valid only when all curvatures and derivatives on space-time background fields are small. Via the σ -model approach one can easily obtain approximate solutions which are in fact identical to the classical equation of gravity in the presence of a dilaton, antisymmetric field and some gauged minimally coupled matter. However, the possible

[†]On leave from Lab. de Physique Théorique de l' ENS, F-75231, Paris, FRANCE.

extension of these approximate solutions to exact string solutions is in general difficult and an unsolved problem at the present time.

The two-dimensional conformal field theory approach takes into account all orders in α' automatically and has the main advantage of providing exact string vacua. The background interpretation of a given exact string solution is a notion which is ill-defined in general. Indeed, the notion of space-time dimensionality and topology breaks down for a solution which involves highly curved backgrounds, namely when the metric and/or gauge field curvatures are of the order of the string scale. A typical example of the space-time dimensional and topological confusion is that of the $SU(2)$ level k group manifold compactification. For large k (small curvature) the target space is a three-dimensional sphere S^3 , $[\frac{SO(4)}{SO(3)} \simeq SO(3) \simeq SU(2)]$. For small k (high curvature) this background interpretation fails. It is in fact well known that the $SU(2)_{k=1}$ WZW model is equivalent to a $c = 1$ conformal system defined by one free bosonic co-ordinate compactified on a cycle with a radius $R = \frac{2}{R} = \sqrt{2}$ (self-dual point).

Naively one may interpret $T_{(R=\sqrt{2})}$ compactification as one-dimensional space with S^1 topology, which is in contradiction to the three-dimensional interpretation with S^3 topology of the $SU(2)_{k=1}$. This shows that both the dimensionality as well as the topology of the target space are not well-defined concepts in string theory. In general, a background interpretation of a given string solution exists only when the lower Kaluza-Klein excitations have masses much smaller than the typical string scale ($M_{st} = \alpha'^{-1/2}$).

In this talk, I will present a class of string solutions which is constructed by using some $N = 4$ superconformal systems with $\hat{c} = 4$ as building blocks. In the limit of small curvatures, these solutions have a non-trivial ten-dimensional background interpretation. Furthermore, the globally defined underlying $N = 4$ worldsheet symmetry stabilizes these solutions under string-loop perturbation, and implies some covariantized space-time supersymmetry around a non-trivial gravitational and dilaton background on the target space.

More explicitly, we arrange the degrees of freedom of the ten supercoordinates in three superconformal systems [2]-[5]:

$$\hat{c} = 10 = \{\hat{c} = 2\} + \{\hat{c} = 4\}_1 + \{\hat{c} = 4\}_2 \quad (1)$$

The $\hat{c} = 2$ subsystem is saturated by two free superfields. In one variation of our solutions, one of the two free superfields is chosen to be the time-like supercoordinate and the other to be one of the nine space-like supercoordinates. In other variations, both supercoordinates are Euclidean or even compactified on a one- or two-dimensional torus.

The remaining eight supercoordinates appear in a group of four in $\{\hat{c} = 4\}_1$ and $\{\hat{c} = 4\}_2$. Both $\{\hat{c} = 4\}_A$ subsystems show an $N = 4$ superconformal symmetry of the

Ademollo et al. type [6]. The non-triviality of our solutions follows from the fact that some realizations of the $\hat{c} = 4, N = 4$ superconformal systems exist which are based on geometrical and topological non-trivial spaces other than the T^4/Z_2 orbifold and the K_3 compact Calabi-Yau space. I will now present two different realizations.

(A) - $W_k^{(4)}$, semi-wormhole realization [7],[8] $\hat{c}[W_k^{(4)}] = 4$

It is based on a supersymmetric version of the $\{U(1) \times SU(2)_k\}$ WZW model. The three bosonic coordinates parametrize the $SU(2)_k$ group manifold while the fourth one is a free field with background charge Q . The four fermionic coordinates are free. In order to obtain a $\hat{c}[U(1) \times SU(2)_k] = 4$ for any value of k , it is necessary to balance the central charge deficit of the $SU(2)_k$ by a central charge benefit of the $U(1)$ background charge Q

$$\begin{aligned}\hat{c}[SU(2)_k] &= \frac{2}{3}\left[3 - \frac{6}{k+2} + \frac{3}{2}\right] = 3 - \frac{4}{k+2} \\ \hat{c}[U(1)_Q] &= \frac{2}{3}\left[1 + 3Q^2 + \frac{1}{2}\right] = 1 + 2Q^2\end{aligned}\tag{2}$$

(The contributions $\frac{3}{2}$ and $\frac{1}{2}$ inside [...] in the first and second line are those of the 3+1 free fermions).

From Eq. (2) one has

$$\hat{c}[SU(2)_k] + \hat{c}[U(1)_Q] = 4 + 2\left(Q^2 - \frac{2}{k+2}\right),$$

and so $\hat{c}[W_k^{(4)}] = 4$ only if

$$Q = \sqrt{\frac{2}{k+2}}\tag{3}$$

The existence of $N = 4, \hat{c} = 4$ superconformal symmetry with this value of Q is found in Ref. [7]. What is extremely interesting is the background interpretation of the $W_k^{(4)}$ space in terms of a four-dimensional (semi)-wormhole space given by Callan, Harvey and Strominger in Ref. [8]. Indeed, for large k , the three $SU(2)_k$ coordinates define a three-dimensional subspace with a non-trivial topology S^3 , while the fourth coordinate with a background charge corresponds to the scale factor of the S^3 sphere.

Another interesting interpretation of the $W_k^{(4)}$ space is its connection with four-dimensional dilaton-axion instantons [8]-[10]. Here, one assumes a string solution based on a six-dimensional compact manifold K^6 with $\hat{c} = 6$; our four-dimensional space-time (after Euclidean rotation) is saturated by $W_k^{(4)}$ with $\hat{c} = 4$.

The explicit realization of the $N = 4$ symmetry of $W_k^{(4)}$, as well as some other properties of string solutions based on it, will be discussed later after I present a new

and inequivalent realization of a $\hat{c} = 4, N = 4$ superconformal system.

B - $\Delta_k^{(4)}$ Cigar-Bell and Trumpet-Bell realizations [5] $\hat{c}[\Delta_k^{(4)}] = 4$

For large values of k (small curvatures), the $\Delta_k^{(4)}$ has a four-dimensional interpretation which is different from $W_k^{(4)}$. The underlying two-dimensional action associated to $\Delta_k^{(4)}$ is an exact superconformal theory based on a supersymmetric version of a gauged WZW model, namely,

$$\Delta_k^{(4)} \equiv \left\{ \left[\frac{SU(2)}{U(1)} \right]_k \times \left[\frac{SL(2, R)}{U(1)} \right]_{k+4} \right\}_{\text{SUSY}} \quad (4)$$

The relation among the levels k and $k' = k + 4$ ensures $\hat{c}[\Delta_k^{(4)}] = 4$ for any k

$$\begin{aligned} \hat{c}\left[\frac{SU(2)}{U(1)}\right]_{ss} &= \frac{2}{3}\left[\frac{3k}{k+2} - 1 + 1\right] = 2 - \frac{4}{k+2} \\ \hat{c}\left[\frac{SL(2, R)}{U(1)}\right]_{ss} &= \frac{2}{3}\left[\frac{3k'}{k'-2} - 1 + 1\right] = 2 + \frac{4}{k'-2} \end{aligned} \quad (5)$$

and so $\hat{c}[\Delta_k^{(4)}] = 4$ when $k' = k + 4$. As we will show later, the above relation among the levels is necessary for the existence of an $N = 4$ superconformal symmetry of Ademollo type [6].

For large k , the $\Delta_k^{(4)}$ space has a dimensional target space interpretation based on a non-trivial background metric G_{ij} and dilaton field Φ given by [11]-[17],[2]-[5]

$$ds^2 = k \left\{ (d\alpha)^2 + \tan^2 \alpha d\theta^2 \right\} + k' \left\{ (d\beta)^2 + \tanh^2 \beta d\varphi^2 \right\} \quad (6a)$$

$$2\Phi = \log \cos^2 \alpha + \log \cosh^2 \beta + \text{const.} \quad (6b)$$

with

$$\alpha \in [0, \frac{\pi}{2}] \cup [\pi, \frac{3}{2}\pi], \quad \beta \in [0, \infty], \quad \theta, \varphi \in [0, 2\pi]$$

The term proportional to k in Eq. (6a) parametrizes the two-dimensional subspace defined by the $\left[\frac{SU(2)}{U(1)} \right]_k$ parafermionic theory [18] and the one proportional to k' is the two-dimensional subspace which is defined by the non-compact parafermionic theory [19],[20],[12] based on the $\left[\frac{SL(2, R)}{U(1)} \right]_{k'}$ axial gauged WZW model. It is well known that a different metric $\tilde{G}_{ij}$ and dilaton function $\tilde{\Phi}$ are obtained if one chooses a vector gauging instead of the axial one [15],[13]:

$$(U(1)^V \times U(1)^V) \rightarrow d\tilde{s}^2 = k \left\{ (d\alpha)^2 + \frac{1}{\tan^2 \alpha} d\theta^2 \right\} + k' \left\{ (d\beta)^2 + \frac{1}{\tanh^2 \beta} d\varphi^2 \right\} \quad (7a)$$

$$2\tilde{\Phi} = \log \sin^2 \alpha + \log \sinh^2 \beta + \text{const.} \quad (7b)$$

In both versions of gaugings, $\Delta_k^{(4)}$ has always one non-compact coordinate (β) and three compact ones (α, θ, φ). (G_{ij}, Φ) and $(\tilde{G}_{ij}, \tilde{\Phi})$ are related by a generalized duality transformation [14]:

$$R(t) \rightarrow \frac{1}{R(t)} , \quad \Phi \rightarrow \Phi + \log R(t) \quad (8)$$

For later purposes, it is more convenient to use the complex notation:

$$z = (\sin \alpha) e^{i\theta} , \quad \omega = (\sinh \beta) e^{i\varphi} , \quad (\text{axial case})$$

$$\text{and} \quad (9)$$

$$\tilde{z} = (\cos \alpha) e^{i\theta} , \quad y = (\cosh \beta) e^{i\varphi} , \quad (\text{vector case})$$

In terms of z and w , G_{ij} and Φ are given by

$$(ds)^2 = k \frac{dz d\bar{z}}{1 - z\bar{z}} + k' \frac{dw d\bar{w}}{w\bar{w} + 1} , \quad (10)$$

and

$$2\Phi = \log(1 - z\bar{z}) + \log(w\bar{w} + 1) + \text{const.}$$

The dual metrics $\tilde{G}_{ij}$ and $\tilde{\Phi}$ are given in terms of $\tilde{z}$ and y as:

$$(d\tilde{s})^2 = k \frac{d\tilde{z} d\tilde{z}^*}{1 - \tilde{z}\tilde{z}^*} + k' \frac{dy d\bar{y}}{y\bar{y} - 1} , \quad (11)$$

and

$$2\tilde{\Phi} = \log(1 - \tilde{z}\tilde{z}^*) + \log(y\bar{y} - 1) .$$

From Eqs. (10) and (11), one observes that the $\left(\frac{SU(2)}{U(1)}\right)_k$ metric and dilaton part are self-dual, (z and $\tilde{z}$), while the $\left(\frac{SU(2,R)}{U(1)}\right)_{k'}$ is not. The w subspace is regular while the y subspace is singular).

The z subspace (or $\tilde{z}$) defines a two-dimensional Bell. Its metric $G_{z\bar{z}}$, the Ricci tensor $R_{z\bar{z}}$ and its scalar curvature $R^{(z)}$ are singular at the boundary of the Bell ($z = 1$):

$$G_{z\bar{z}} = \frac{k}{1 - z\bar{z}} , \quad R_{z\bar{z}} = \frac{-1}{(1 - z\bar{z})^2} , \quad R^{(z)} = \frac{-1}{k(1 - z\bar{z})} , \quad (12)$$

The z -Bell is negatively curved for any value of $|z| < 1$.

The w subspace is regular everywhere with finite positive curvature and is asymptotically flat for $|w| \rightarrow \infty$

$$G_{w\bar{w}} = \frac{k'}{w\bar{w} + 1} , \quad R_{w\bar{w}} = \frac{+1}{(w\bar{w} + 1)^2} , \quad R^{(w)} = \frac{+1}{k'(w\bar{w} + 1)} , \quad (13)$$

It has a cigar shape with maximal curvature at $w = 0$.

The y subspace has a different shape from its w -dual. It looks like a two-dimensional trumpet with infinite curvature at the boundary ($y = 1$). It is positively curved everywhere ($|y| > 1$) and is asymptotically flat for $|y| \rightarrow \infty$:

$$G_{y\bar{y}} = \frac{k'}{y\bar{y} - 1} , \quad R_{y\bar{y}} = \frac{+1}{(y\bar{y} - 1)^2} , \quad R^{(y)} = \frac{+1}{k'(y\bar{y} - 1)} , \quad (14)$$

We have therefore two different versions of the $\Delta_k^{(4)}$ space. The first one is that of the (z, w) -BELL-CIGAR four-dimensional space and the second one is that of the $(\tilde{z}, y)$ BELL-TRUMPET four-dimensional space.

Up to now we have discussed the geometrical structure of the bosonic coordinates of the $\Delta_k^{(4)}$ space. In order to complete the description of our solution, we must include the fermionic superpartners of (z, w) coordinates, e.g., four Weyl Majorana left-handed two-dimensional fermions ($\psi_+^a, a = 1, 2, 3, 4$) as well as four Weyl-Majorana right-handed ones ($\psi_-^a, a = 1, 2, 3, 4$).

Because of the $N = 1$ local supersymmetry, the interactions among fermions are fixed in terms of the two-dimensional σ -model backgrounds G_{ij}, B_{ij} and Φ). In the superconformal gauge, one has the following generic form for the $N = 1$ σ -model action (in the absence of $B_{ij} = 0$):

$$S = -\frac{1}{2\pi} \int d\xi d\bar{\xi} \left\{ V_+^a V_-^a - \frac{1}{2} (\psi_+^a \nabla_- \psi_+^a - \psi_-^a \nabla_+ \psi_-^a) - \frac{1}{2} R_{ab,cd} \psi_+^a \psi_+^b \psi_-^c \psi_-^d + \Phi R^{(2)} \right\} \quad (15)$$

where $a = 1, 2, 3, 4$ are local flat indices and

$$V_+^a = E_i^a \partial X^i , \quad V_-^a = E_i^a \bar{\partial} X^i , \quad \text{with} \quad G_{ij} = E_i^a E_j^a \quad (16)$$

The ∇_- and ∇_+ denote the left- and right-handed covariant derivatives acting on left- and right-handed fermions ψ_+^a, ψ_-^a :

$$\begin{aligned} \nabla_- \psi_+^a &= \bar{\partial} \psi_+^a + \Gamma_{bc}^a \psi_+^c V_-^b \\ \nabla_+ \psi_-^a &= \partial \psi_-^a + \Gamma_{bc}^a \psi_-^c V_+^b \end{aligned} \quad (17)$$

Since $\Delta_K^{(4)}$ is defined as a direct product of two two-dimensional subspaces, the σ -model action can be written as:

$$S(\Delta_K^{(4)}) = S \left[\frac{SU(2)}{U(1)} \right]_k + S \left[\frac{SL(2, R)}{U(1)} \right]_{k'} \quad (18)$$

Because of the identity

$$R_{ab,cd} \psi_+^a \psi_+^b \psi_-^c \psi_-^d = 2R \psi_+^1 \psi_+^2 \psi_-^1 \psi_-^2 , \quad (19)$$

valid in both subspaces and thanks to the relations [5]

$$-(R^{(z)} + \Gamma_z G^{z\bar{z}} \Gamma_{\bar{z}}) = \frac{1}{k} ; \left(\frac{SU(2)}{U(1)} \right)_k \quad (20a)$$

$$-(R^{(w)} + \Gamma_w G^{z\bar{z}} \Gamma_{\bar{w}}) = \frac{-1}{k'} ; \left(\frac{SL(2,R)}{U(1)} \right)_{k'} \quad (20b)$$

it is possible to rewrite the σ -model action in a more convenient form which shows in particular (at least at the classical level) that the fermions can be described in terms of two free bosonic fields compactified in a special radius. Indeed, using Eqs. (19) and (20), one finds [5]:

$$\begin{aligned} S \left[\frac{SU(2)}{U(1)} \right]_k &= \frac{-1}{4\pi} \int d\xi d\bar{\xi} \left\{ \partial A \bar{\partial} A + t(A)^2 (\partial \theta_A - \sqrt{\frac{2}{k}} \psi_+^1 \psi_+^2) (\bar{\partial} \theta_A - \sqrt{\frac{2}{k}} \psi_-^1 \psi_-^2) \right. \\ &\quad - (\psi_+^1 \bar{\partial} \psi_+^2 + \psi_+^2 \bar{\partial} \psi_+^1 + \psi_-^1 \partial \psi_-^1 + \psi_-^2 \partial \psi_-^2) \\ &\quad \left. + \frac{2}{k} (\psi_+^1 \psi_+^2) (\psi_-^1 \psi_-^2) + \log C^2(A) R^{(2)} \right\} \end{aligned} \quad (21)$$

where A, θ_A are rescaled fields so that, in the large k limit, real bosonic fields are conventionally normalized [see Eq. (9)]:

$$\begin{aligned} \sin^2 \frac{A}{\sqrt{2k}} &= \sin^2 \alpha = z\bar{z} \\ -i \frac{1}{2} \log \frac{z}{\bar{z}} &= \theta = \frac{\theta_A}{\sqrt{2k}} \\ t(A)^2 &= \frac{z\bar{z}}{1 - z\bar{z}} = \tan^2 \frac{A}{\sqrt{2k}} \\ C(A)^2 &= 1 - z\bar{z} = \cos^2 \frac{A}{\sqrt{2k}} \end{aligned} \quad (22)$$

In a similar way, the

$$\begin{aligned} S \left[\frac{SL(2,R)}{U(1)} \right]_{k'} &= \frac{-1}{4\pi} \int d\xi d\bar{\xi} \left\{ \partial B \bar{\partial} B + T(B)^2 (\partial \theta_B - \sqrt{\frac{2}{k'}} \psi_+^3 \psi_+^4) (\bar{\partial} \theta_B - \sqrt{\frac{2}{k'}} \psi_-^3 \psi_-^4) \right. \\ &\quad - (\psi_+^3 \bar{\partial} \psi_+^4 + \psi_+^4 \bar{\partial} \psi_+^3 + \psi_-^3 \partial \psi_-^3 + \psi_-^4 \partial \psi_-^4) \\ &\quad \left. - \frac{2}{k'} (\psi_+^3 \psi_+^4) (\psi_-^3 \psi_-^4) + \log C(B)^2 R^{(2)} \right\} \end{aligned} \quad (23)$$

where B and θ_B are now defined in terms of w fields:

$$\begin{aligned} \sinh^2 \frac{B}{\sqrt{2k'}} &= \sinh^2 \beta = w\bar{w} \\ -i \frac{1}{2} \log \frac{w}{\bar{w}} &= \varphi = \frac{\theta_B}{\sqrt{2k'}} \\ T(B)^2 &= \frac{w\bar{w}}{1 + w\bar{w}} = \tanh^2 \frac{B}{\sqrt{2k'}} \\ C(B)^2 &= 1 + w\bar{w} = \cosh^2 \frac{B}{\sqrt{2k'}} \end{aligned} \quad (24)$$

In both Eqs. (21) and (23), the pure fermionic part of the action is given by the free-fermion kinetic terms together with some current-current interaction. This fact permits us to describe the four left and the four right fermions in terms of free bosonic fields ϕ_A, ϕ_B compactified in the shifted radii [5]:

$$R_A^2 = 1 + \frac{2}{k} = \frac{k+2}{k} ; \quad \left(\frac{SU(2)}{U(1)} \right)_k \quad (25a)$$

$$R_B^2 = 1 - \frac{2}{k'} = \frac{k'-2}{k'} ; \quad \left(\frac{SL(2,R)}{U(1)} \right)_{k'} \quad (25b)$$

The deviation from the value $R_A = R_B = 1$ is due to current-current interactions. The decoupling of ϕ_A and ϕ_B fields can be seen via the bosonization of fermions and the redefinition of the $\overset{(-)}{\partial} \theta_A$ and $\overset{(-)}{\partial} \theta_B$ bosonic currents order by order in a $\frac{1}{k}$ or $\frac{1}{k'}$ expansion. This statement is indeed exact and it follows from the fact that both $\left(\frac{SU(2)}{U(1)} \right)_k$ and $\left(\frac{SL(2,R)}{U(1)} \right)_{k'}$ are exact $N = (2, 2)$ superconformal models. This fact is well known in both $\frac{SU(2)}{U(1)}$ and $\frac{SL(2,R)}{U(1)}$ supersymmetric coset models in the compact and non-compact parafermionic representations. The $N = 2$ generators, $J(\xi), G(\xi), \bar{G}(\xi)$ and $T(\xi)$ are given in terms of free-scalar fields (ϕ_A, ϕ_B) and in terms of (non-local) parafermionic currents [18]-[21], $(P_k, \Pi_{k'})$,

$$(i) - \left(\frac{SU(2)}{U(1)} \right)_k$$

$$\begin{aligned} J(\xi) &= \sqrt{\frac{k}{k+2}} \partial \phi_A \\ \overset{(-)}{G}(\xi) &= \overset{(-)}{P}_k e^{\pm i \sqrt{\frac{k+2}{k}} \phi_A} \\ T(\xi) &= -\frac{1}{2} (\partial \phi_A)^2 + T_P(\xi) \end{aligned} \quad (26)$$

$$(ii) - \left(\frac{SL(2,R)}{U(1)} \right)_{k'}$$

$$\begin{aligned} J(\xi) &= \sqrt{\frac{k'}{k'-2}} \partial \phi_B \\ \overset{(-)}{G}(\xi) &= \overset{(-)}{\Pi}_{k'} e^{\pm i \sqrt{\frac{k'-2}{k'}} \phi_B} \\ T(\xi) &= -\frac{1}{2} (\partial \phi_B)^2 + T_\Pi(\xi) \end{aligned} \quad (27)$$

ϕ_A and ϕ_B are free bosons compactified on radii $R_A = \sqrt{\frac{k+2}{k}}$ and $R_B = \sqrt{\frac{k'-2}{k'}}$ and parametrize the fermions $\psi_\pm^a, a = 1, 2, 3, 4$ which appear in the σ -model actions in Eqs. (21) and (23) according to our previous discussion. For large k, k' , the operators $e^{i \sqrt{\frac{k+2}{k}} \phi_A}$

and $e^{i\sqrt{\frac{k'-2}{k'}}\phi_B}$ have conformal dimensions of almost $\frac{1}{2}$ ($h_A = \frac{1}{2} + \frac{1}{k}$, $h_B = \frac{1}{2} - \frac{1}{k'}$) P_k and $\Pi_{k'}$ are the parafermionic currents with dimensions $h_P = 1 - \frac{1}{k}$ and $h_\Pi = 1 + \frac{1}{k'}$, so that $G(\xi)$ in Eqs. (26) and (27) are the $N = 2$ supercurrents of dimension $\frac{3}{2}$.

As usual, the compact and non-compact parafermions satisfy the algebra which follows from their O.P.E. [18]-[20]:

$$\begin{aligned} P_k(\xi)\bar{P}_k(\xi') &= 2\left[\left(\frac{k}{k+2}\right)\frac{1}{(\xi-\xi')^2} + T_P(\xi')\right](\xi-\xi')^{\frac{2}{k}} \\ \Pi_{k'}(\xi)\bar{\Pi}_{k'}(\xi') &= 2\left[\left(\frac{k'}{k'-2}\right)\frac{1}{(\xi-\xi')^2} + T_\Pi(\xi')\right](\xi-\xi')^{-\frac{2}{k'}} \end{aligned} \quad (28)$$

and the dimension-2 operators $T_P(\xi)$ and $T_\Pi(\xi')$ are their corresponding stress tensors. $T_P(\xi)$ and $T_\Pi(\xi')$ satisfy the Virasoro algebra

$$T(\xi)T(\xi') = \frac{c}{2(\xi-\xi')^4} + \frac{2T(\xi')}{(\xi-\xi')^2} + \frac{\partial T(\xi')}{(\xi-\xi')} \quad (29)$$

with central charge $c_P = \frac{3k}{k+2} - 1$, $c_\Pi = \frac{3k'}{k'-2} - 1$. Using the above O.P.E. [Eqs. (28), (29)] and those of the free fields

$$\partial\phi_A \partial\phi_A = -\frac{1}{(\xi-\xi')^2}, \quad \partial\phi_B \partial\phi_B = -\frac{1}{(\xi-\xi')^2}, \quad (30)$$

it is easy to show the closure of the $N = 2$ algebra for both $\left(\frac{SU(2)}{U(1)}\right)_k$ and $\left(\frac{SL(2,R)}{U(1)}\right)_{k'}$ supersymmetric coset models.

What is interesting is the interpretation of the almost dimension-one, non-local cite-bkk currents P_k and $P_{i_{k'}}$ for k and k' large citeee. P_k are the conjugate momenta of the two-dimensional subspace z while $P_{i_{k'}}$ are those of the w subspace. As we mention in the introduction, this interpretation breaks down for small values of k and k' , e.g., when the target backgrounds are strongly curved.

For large values of k and k' , P_k generalizes two out of the four flat space currents while $\Pi_{k'}$ generalizes the two remaining ones [5], [23]

$$\begin{aligned} \overset{(-)}{P}_k &\xrightarrow{k \rightarrow \infty} \pm \partial x_1 + i \partial x_2 \\ \overset{(-)}{\Pi}_{k'} &\xrightarrow{k' \rightarrow \infty} \pm \partial x_4 + i \partial x_3 \end{aligned} \quad (31)$$

We would like to show that for $k' = k + 4$ the $N = 2$ superconformal symmetry is extended to $N = 4$ with $\hat{c} = 4(c = 6)$ for any value of k . For this purpose it is more convenient to redefine the fields ϕ_A and ϕ_B in terms of two other bosonic fields H_+ and H_- as follows:

$$\sqrt{2}H_+ = \sqrt{\frac{k}{k+2}} \phi_A + \sqrt{\frac{k+4}{k+2}} \phi_B$$

$$\sqrt{2}H_- = +\sqrt{\frac{k+4}{k+2}} \phi_A - \sqrt{\frac{k}{k+2}} \phi_B \quad (32)$$

The $N = 2$ current of the $\Delta_k^{(4)}$ is then given uniquely in terms of the ∂H_+ current. It is then possible (due to the $N = 2$ algebra) to factor the H_+ dependence from the (complex) supercurrent. One finds [5]

$$G = \left[P_k e^{i\frac{\alpha}{\sqrt{2}}H_-} + \Pi_{k'} e^{-i\frac{1}{\alpha\sqrt{2}}H_-} \right] e^{i\frac{1}{\sqrt{2}}H_+} \quad (33)$$

with $\alpha = \sqrt{\frac{k+4}{k}}$.

It is suggestive to compare the above expression of the supercurrent with that of the free-field realization [6] ($k \rightarrow \infty, \alpha = 1$)

$$G_{free} = \left[(\partial x_1 + i\partial x_2) e^{i\frac{1}{\sqrt{2}}H_-} + (\partial x_4 + i\partial x_3) e^{-i\frac{1}{\sqrt{2}}H_-} \right] e^{i\frac{1}{\sqrt{2}}H_+} \quad (34)$$

Here H_+ and H_- are given in terms of free fermions via the conventional bosonization

$$\begin{aligned} \sqrt{2}H_+ &= \psi_1\psi_2 + \psi_4\psi_3 \\ \sqrt{2}H_- &= \psi_1\psi_2 - \psi_4\psi_3 \end{aligned} \quad (35)$$

There are some similarities and some basic differences between the two realizations. First, the rôle of the H_+ field is the same and, as is expected from the $N = 2$ algebra, is a free field compactified on a radius $R_{H^+} = \sqrt{\frac{c}{3}} = \sqrt{2}$. This value of the radius is special and extends in both cases the $U(1)$ to $SU(2)_1$ current algebra. In free field realization, however, the $N = 2$ superconformal symmetry is extended to an $N = 4$, due to the existence of an additional supercurrent, namely:

$$\tilde{G}_{free} = \left[(\partial x_1 - i\partial x_2) e^{-i\frac{1}{\sqrt{2}}H_-} - (\partial x_4 - i\partial x_3) e^{i\frac{1}{\sqrt{2}}H_-} \right] e^{i\frac{1}{\sqrt{2}}H_+}$$

In a similar way, an additional supercurrent exists also in the $\Delta_k^{(4)}$ realization [5]:

$$\tilde{G} = \left[\bar{P}_k e^{-i\frac{\alpha}{\sqrt{2}}H_-} - \bar{\Pi}_{k'} e^{i\frac{1}{\alpha\sqrt{2}}H_-} \right] e^{i\frac{1}{\sqrt{2}}H_+} \quad (36)$$

The $N = 4$ superconformal algebra closes in both realizations among $[T_B, G, \tilde{G}$ and $S_i]$ with central charge $\hat{c} = 4$. This shows that the $\Delta_k^{(4)}$ space shares the same global superconformal symmetries as the T^4/Z_2 orbifold model, the four-dimensional Calabi-Yau space K_3 and the previously discussed $W_k^{(4)}$ wormhole space. $\Delta_k^{(4)}$ and $W_k^{(4)}$ are not only non-trivial spaces but are also non-compact spaces (K_3 and T^4/Z_2 are compact).

For completeness, we will present below the basic operators and fields of the $W_k^{(4)}$ realization [7],[8],[2]-[5],[10]. As we already mentioned, $W_k^{(4)}$ is based on a supersymmetric

$SU(2)_k \times U(1)_Q$ WZW model with a background term $Q = \sqrt{\frac{2}{k+2}}$ in the $U(1)_Q$ current. The four fermions of the model are free and are parametrized by the H^+ and H^- fields (via bosonization) as in the free-field representation [Eq. (35)]. The four coordinate currents are the three $SU(2)_k$ ($J^i, i = 1, 2, 3$) currents and the $U(1)_Q$ J_4 current

$$\begin{aligned} J^i(\xi)J^i(\xi') &= -\frac{k}{2} \frac{\delta^{ij}}{(\xi - \xi')^2} + \epsilon^{ij\ell} \frac{J^\ell}{(\xi - \xi')^2}, \quad i = 1, 2, 3 \\ J^4(\xi)J^4(\xi') &= \frac{-1}{(\xi - \xi')^2} \cdot \frac{1}{Q^2} \end{aligned} \quad (37)$$

(the Q rescaling of J^4 is for convenience).

The $T_B, B, \tilde{G}$ and S_i associated to $W_k^{(4)}$ are

$$\begin{aligned} T_B &= -\frac{1}{2} \left[(\partial H^+)^2 + (\partial H^-)^2 + Q^2 (J_1^2 + J_2^2 + J_3^2 + J_4^2 + \partial J_4) \right] \\ G &= Q \left[(J_1 + iJ_2) e^{i\frac{1}{\sqrt{2}}H^-} + (J_4 + i(J_3 + \sqrt{2}\partial H^-)) e^{i\frac{1}{\sqrt{2}}H^-} \right] e^{i\frac{1}{\sqrt{2}}H^+} \\ \tilde{G} &= Q \left[(J_1 - iJ_2) e^{i\frac{1}{\sqrt{2}}H^-} - (J_4 - i(J_3 + \sqrt{2}\partial H^-)) e^{i\frac{1}{\sqrt{2}}H^-} \right] e^{i\frac{1}{\sqrt{2}}H^+} \\ S_0 &= \frac{1}{\sqrt{2}}\partial H^+, \quad S_\pm = e^{\pm i\sqrt{2}H^\pm} \end{aligned} \quad (38)$$

In the $W_k^{(4)}$ realization, both H^+ and H^- are compactified on a torus with radius $R_{H^+} = R_{H^-} = \sqrt{2}$ as in the free field case. There are in total three underlying $SU(2)$ Kac-Moody currents: (i) the $SU(2)_k$ defined by the coordinate currents, (ii) the $SU(2)_1^+$ defined by the H^+ field $R_{H^+} = \sqrt{2}$, (iii) the $SU(2)_1^-$ defined by the H^- field $R_{H^-} = \sqrt{2}$.

The background term in $T_B, (Q\partial J^4)$, comes from a non-trivial dilaton background: $\Phi = QX^4$. The term $Q\sqrt{2}\partial H^- \exp[\pm i\frac{1}{\sqrt{2}}H^- + i\frac{1}{\sqrt{2}}H^+]$ in G and $\tilde{G}$ describes, at the same time, the standard fermionic torsion term $\pm Q\psi^i\psi^j\psi^k$ ($i = 1, 2, 3, 4$) as well as the fermionic background term $\pm Q\partial\psi^i$. They are arranged together in $\sqrt{2}\partial H^-$ which shifts the J_3 current. Contrary to the $\Delta_k^{(4)}$ space in which the torsion terms were absent, in $W_k^{(4)}$ space there is a non-vanishing torsion due to the WZ term proportional to the $SU(2)$ structure constant. So, in $W_k^{(4)}$ the antisymmetric field background B_{ij} is non-zero with non-trivial field strength H_{ijk}

$$H_{ijk} = e^{-2\phi} \epsilon_{ijk}{}^\ell \partial_\ell \Phi \simeq e^{-2\phi} Q \cdot \epsilon_{ijk}{}^4 \quad (39)$$

Summarizing our results, we propose a special class of exact string solutions [2]-[5] with $N = (1, 0)$ or $N = (1, 1)$ local worldsheet supersymmetry. In the type II case, we arrange the degrees of freedom of the ten supercoordinates in three superconformal systems:

$$\hat{c} = 10 = \{\hat{c} = 2\}_0 + \{\hat{c} = 4\}_1 + \{\hat{c} = 4\}_2$$

The $\hat{c} = 2$ system is saturated by two free superfields (compact or non-compact), and so the background metric is flat:

$$ds^2\{F^2\} = dx d\bar{x} \ , \ (x = x_1 + ix_2) \ .$$

The remaining eight supercoordinates appear in a group of four. In both $\{\hat{c} = 4\}_{1,2}$ subsystems an $N = 4$ globally defined superconformal symmetry is assumed. We propose three exact $N = 4$ superconformal systems with $\hat{c} = 4$.

(i) - *Flat realization* : $\hat{c} = 4$

$$ds^2(F) = dz d\bar{z} + dw d\bar{w}$$

$$\Phi = \text{constant} \ , \ B_{ij} = \text{constant}$$

This realization includes the flat space as well as the toroidal T^4 and orbifold T^4/Z_2 models.

(ii) - *Semi – wormhole space* $W_k^{(4)}$ with $\hat{c} = 4$

$$ds^2(W) = k \frac{dz d\bar{z} + dw d\bar{w}}{z\bar{z} + w\bar{w}}$$

$$2\Phi = \log(z\bar{z} + w\bar{w})$$

$$H_{ijk} = \epsilon_{ijkl} \partial^\ell \Phi$$

(iii) - *The $\Delta_k^{(4)}$ realization* with $\hat{c} = 4$

$$ds^2(\Delta) = k \frac{dz d\bar{z}}{1 - z\bar{z}} + (k + 4) \frac{dw d\bar{w}}{w\bar{w} + \epsilon}$$

$$2\Phi(\Delta) = \log(1 - z\bar{z}) + \log(w\bar{w} + \epsilon) \ , \ \epsilon = \pm 1$$

$$H_{ijk} = 0$$

$$\epsilon = 1 \ , \ \text{Bell – Cigar space}$$

$$\epsilon = -1 \ , \ \text{Bell – Trumpet space}$$

Using as building blocks the $F^2, F^4, \Delta_\epsilon^4$ or W^4 superconformal systems, one obtains several string solutions based on non-trivial ten-dimensional backgrounds. The advantage of our solutions compared to the subclass of models $(F^6 \times W^{(4)})$ studied in the literature [8] lies in the complete knowledge of the $\{\hat{c} = 2\}_0$ and $\{\hat{c} = 4\}_A$ superconformal theories.

Due to this we are able to study not only the background solutions for large k_A but also the full string spectrum and derive the partition functions of the models in a modular invariant way respecting the $N = 4$ superconformal symmetry [2]-[5]. Indeed, the full spectrum is given in terms of several characters of known conformal theories. In the W_k^4 realization [2]-[4] one uses a character combination of $SU(2)_k$, $SU(2)_1^+$ and $SU(2)_1^-$ together with the $U(1)_Q$ Liouville-type characters. In $\Delta_k^{(4)}$, one [5] uses the compact [24] and non-compact [25] parafermionic characters (string functions) together with the $SU(2)_1^+$ and $U(1)_{R_H^-}$ ($R_H = \sqrt{2} \sqrt{\frac{k+4}{k}}$). Finally in F^2 and F^4 , one uses non-compact, toroidal or orbifold characters.

It is important to stress here that the character combinations above, are not arbitrary but are dictated by the global existence of $N = 4$ superconformal symmetry as well as modular invariance.

These requirements define some generalized GSO projections [2]-[5], similar to that of the fermionic construction [26] and that of the conformal block construction [27] of Gepner. One of these projections is fundamental and guarantees the existence of some space-time supersymmetries via the $N = 4$ spectral flows. It can be expressed in terms of the two $N = 4$ ($\hat{c} = 4$) $_A$, $A = 1, 2$ $\{SU(2)_1^+\}_A$ spins, $[j(s_1), j(s_2)]$ [2]-[5]:

$$2(j(s_1) + j(s_2)) = \text{odd integer} . \quad (40)$$

Equation (40) guarantees the existence of some covariantized target space symmetries around the proposed non-trivial backgrounds. This stabilizes our solutions (at least perturbatively) and projects out all kinds of tachyonic or complex conformal weight states. This projection phenomenon is similar to the one observed by Kutasov and Seiberg in the framework of non-critical superstrings with an $N = 2$ globally defined superconformal symmetry [28].

Heterotic solutions are simply obtained via a generalized [29],[27] heterotic map. Here also the solutions are stable but the number of covariantized space-time supersymmetries is reduced by a factor of two.

We hope that our explicit construction of a family of consistent and stable solutions will give a better understanding of some fundamental string properties, especially in the case of strongly curved backgrounds (small k_A) where the notion of the space-time dimensionality and topology breaks down.

ACKNOWLEDGEMENTS

I would like to thank I. Antoniadis and S. Ferrara. Some results about the $W_k^{(4)}$ space of this work are products of our collaboration. I am also grateful to C. Callan, D. Gross, E. Kiritsis and D. Lüst for fruitful discussions.

References

- [1] C. Callan, D. Friedman, E. Martinec and M. Perry, *Nucl.Phys.* **B262** (1985) 593;
E. Fradkin and A. Tseytlin, *Nucl.Phys.* **B261** (1985) 1.
- [2] I. Antoniadis, S. Ferrara and C. Kounnas, LPTENS-CERN preprint in preparation.
- [3] I. Antoniadis and C. Kounnas, LPTENS-CERN preprint in preparation.
- [4] C. Kounnas, Proceedings of the Geneva International Conference, on “Lepton-photon Symposium and Europhysics Conference on High Energy Physics”, 25 July - 1 August 1991, Vol. 1, pp. 302-306.
- [5] C. Kounnas, CERN preprint, CERN-TH.6799/93.
- [6] M. Ademollo et al., *Nucl.Phys.* **B114** (1976) 297;
T. Eguchi and A. Taormina, *Phys.Lett.* **B200** (1988) 634;
A. Sevrin, W. Troost and A. Van Proeyen, *Phys.Lett.* **B208** (1988) 447.
- [7] C. Kounnas, M. Porrati and B. Rostand, *Phys.Lett.* **B258** (1991) 61.
- [8] C. Callan, J. Harvey and A. Strominger, *Nucl.Phys.* **B359** (1991) 611;
C. Callan, Lectures at Sixth J.A. Swieca Summer School, PUPT-1278 (1991).
- [9] S.J. Rey, *Phys.Rev.* **D43** (1991) 526 and SLAC-PUB-5659 (1991).
- [10] M. Billó, P. Fré, L. Girardello and A. Zaffaroni, preprint SISSA 159/92/EP or IFUM/431/FT.
- [11] E. Witten, *Phys.Rev.* **44** (1991) 314.
- [12] I. Bars, *Nucl.Phys.* **B334** (1990) 125;
I. Bars and D. Nemeschansky, *Nucl.Phys.* **B348** (1991) 89.
- [13] R. Dijkraaf, E. Verlinde and H. Verlinde, *Nucl.Phys.* **B371** (1992) 269.

- [14] A. Giveon, *Mod.Phys.Lett.* **A6** (1991) 2843.
- [15] E. Kiritsis, *Mod.Phys.Lett.* **A6** (1991) 2871; CERN preprint, CERN-TH.6797/93.
- [16] P. Ginsparg and F. Quevedo, *Nucl. Phys.* **B385** (1992) 527.
- [17] C. Kounnas and D. Lüst, *Phys.Lett.* **B289** (1992) 56.
- [18] A.B. Zamolodchikov and V.A. Fateev, *Sov.Phys.-JETP* **62** (1985) 215.
- [19] L. Dixon, J. Lykken and M. Peskin, *Nucl.Phys.* **B325** (1989) 329.
- [20] J. Lykken, *Nucl.Phys.* **B313** (1989) 473.
- [21] A.B. Zamolodchikov and V.A. Fateev, *Sov.Phys.-JETP* **63** (1986) 913;
P. Di Vecchia, J. Petersen, M.Yu and H. Zheng, *Phys.Lett.* **B174** (1986) 280;
Z. Qiu, *Phys.Lett.* **B188** (1987) 207.
- [22] I. Bakas and E. Kiritsis, ENS preprint, LPTENS-92-30 (1992).
- [23] I. Bakas, E. Kiritsis, *Nucl. Phys.* **B343** (1990) 185; *Mod. Phys. Lett.* **A5** (1990) 2039.
- [24] V. Kac and D. Peterson, *Adv.Math.* **53** (1984) 125.i
- [25] I. Bakas and E. Kiritsis, *Int.J.Mod.Phys.* **A7** [Suppl. 1A] (1992) 339.
- [26] I. Antoniadis, C. Bachas and C. Kounnas, *Nucl.Phys.* **B289** (1987) 87;
H. Kawai, D.C. Lewellen and S.H.-H. Tye, *Nucl.Phys.* **B288** (1987) 1.
- [27] D. Gepner, *Phys.Lett.* **B199** (1987) 370; *Nucl.Phys.* **B296** (1988) 757.
- [28] D. Kutasov and N. Seiberg, *Phys.Lett.* **B251** (1990) 67.
- [29] W. Lerche, D. Lüst and A.N. Schellekens, *Phys.Lett.* **B181** (1986) 71;
W. Lerche, D. Lüst and A.N. Schellekens, *Nucl.Phys.* **B287** (1987) 477.